



## Personality factors and eating disorders: Self-uncertainty

Astrid von Lojewski <sup>a,\*</sup>, Suzanne Abraham <sup>a,b</sup>

<sup>a</sup> Department of Obstetrics and Gynaecology, University of Sydney, Royal North Shore Hospital, Building 52, Pacific Highway, St Leonards, NSW 2065, Australia

<sup>b</sup> The Northside Clinic, 2 Greenwich Road, Greenwich, NSW 2065, Australia



### ARTICLE INFO

#### Article history:

Received 13 April 2013

Received in revised form 22 September 2013

Accepted 23 October 2013

Available online 31 October 2013

#### Keywords:

Eating disorder

Factor analysis

Personality

Self-concept

Perfectionism

Avoidance

### ABSTRACT

The International Personality Disorder Examination interview (IPDE) was used to examine common features of personality amongst eating disorder (ED) patients. Female inpatients (N = 155), aged 18 to 45, BMI < 30 kg/m<sup>2</sup>, were interviewed. Items present in ≥25% of patients were analysed by factor analysis. Five factors emerged — ‘interpersonal anxiety’, ‘instability’, ‘self-uncertainty’, ‘obsessionality’ and ‘perfectionism’ accounting for 62% of the variance. Patients with BMI, <18.5 kg/m<sup>2</sup> had significantly greater ‘interpersonal anxiety’ factor scores. Patients who purged had higher ‘interpersonal anxiety’, ‘instability’, and ‘perfectionism’ factor scores. Differences between ED diagnostic groups were accounted for by body weight and purging. Increasing age was weakly associated with improvement in ‘self-uncertainty’ and ‘instability’ scores. This study separates obsessionality and perfectionism, possibly reflecting ED patients’ ‘need for control’, and introduces a new factor ‘self-uncertainty’ which reflects their poor self-concept. The contribution of this factor structure to development and duration of illness should be studied.

© 2013 Elsevier Ltd. All rights reserved.

### 1. Introduction

Both presence and severity of personality pathology influence the course of psychiatric disorders (Grilo, 2002). Personality features are thought to be relevant to general functioning, clinical characteristics and prognosis of persons with eating disorders (ED); greater pathology requires more intense treatment (Klump et al., 2004; Larsson & Hellzen, 2004; Lilienfeld, Wonderlich, Riso, Crosby, & Mitchell, 2006; Milos, Spindler, Buddeberg, & Cramer, 2003; Steiger & Bruce, 2004; Treasure, Crane, McKnight, Buchanan, & Wolfe, 2011; Wagner et al., 2006; Westen & Harnden-Fischer, 2001).

Research into personality disorders (PD) in ED has yielded inconsistent results (Diaz-Marsa, Carrasco Perera, Prieto Lopez, & Saiz Ruiz, 2000; Godt, 2008; Ilkjaer et al., 2004; Kennedy, McVey, & Katz, 1990; Modestin, Oberson, & Erni, 1997; Rosenvinge, Martinussen, & Ostensen, 2000; Vitousek & Manke, 1994; Zubieta, Demitrack, Fenick, & Krahn, 1995) and prevalence rates of PD in ED vary — between 27% and 93% (Skodol et al., 1993). Contributing to these variations are: 1. use of different interview schedules (Skodol et al., 1993; Wonderlich et al., 2007) and measurement tools such as self-report instruments, which overestimate the prevalence of personality pathology (Cassin & von Ranson, 2005); 2. assessment of personality as categorical or dimensional; the former having poorer discriminant validity (Haslem, Holland, & Kuppens, 2012; Loranger et al., 1994; Wonderlich & Mitchell, 2001); 3. temporal instability of PD categories (Wonderlich & Mitchell, 2001) and 4. ED behaviours may influence reporting of personality variables. Additionally,

previous factor analytical studies of personality pathology in ED have identified varying factor structures often differing from existing personality constructs (Grilo, 2004; Lampard, Byrne, McLean, & Fursland, 2012). This suggests the importance of exploring personality factors in ED.

The International Personality Disorder Examination (IPDE; Loranger, Janca, & Sartorius, 1997) is considered an accurate, conservative instrument and ‘better than clinical judgment’ (Karwautz, Troop, Rabe-Hesketh, Collier, & Treasure, 2003). Providing dimensional scores for DSM-IV PD, it has been validated in a worldwide WHO study revealing good inter-rater reliability and temporal stability (Loranger et al., 1994). The interview contains prompt questions and asks for examples to clarify and ensure that the item is not attributable to the ED. Previously the IPDE has only been used to investigate differences between obese, binge eating disorder (BED) and bulimia nervosa (BN) groups (van Hanswijck de Jonge, van Furth, Lacey, & Waller, 2003).

We explored the factor structure derived from the more common and frequently cited set of items of the IPDE within a non-obese ED population (trans-diagnostic approach), and ascertained if these differed or were common for patients: employing different weight losing behaviours, at different body weights and, post hoc, with different diagnoses.

### 2. Material and methods

#### 2.1. Participants

Patients consecutively admitted to the Northside Clinic ED unit for inpatient treatment were asked to participate if they were female, 18 to 45 years, and had no major comorbid psychiatric (e.g. bipolar disorder) or medical diagnoses (e.g. diabetes). All patients fulfilled the

\* Corresponding author. Tel.: +61 2 9926 8308; fax: +61 2 9906 6742.

E-mail addresses: astrid.vonlojewski@sydney.edu.au (A. von Lojewski), suzanne.abraham@sydney.edu.au (S. Abraham).

DSM-IV criteria for a current ED (American Psychiatric Association (APA), 1994).

Of the 171 eligible patients approached 16 did not give consent. There were no significant differences between those participating or not participating for age and current BMI. Of the 155 participants included in the study 34 had a diagnosis of anorexia nervosa-restricting type (AN-R), 29 a diagnosis of anorexia nervosa-binge/purging type (AN-BP), 44 had BN and 48 were diagnosed with eating disorder not otherwise specified (EDNOS). Average age of sample was 24 years (SD 6) and there was no significant difference between the ED diagnostic groups on ED QOL Global Score (mean 16, SD 4).

2.2. Measures

The IPDE is a semi-structured clinical diagnostic interview created as a standardised and international resource to assess PDs. The items cover all criteria for the ten Axis II disorder diagnoses of the DSM-IV. All items are rated as definite (score 2), probable (score 1) or absent (score 0). The dimensional score is calculated by adding up all item scores of a PD (0, 1 or 2). The dimensional scores were used as they have higher reliabilities than definite categorical scores (score 2) (Loranger et al., 1994).

Additional clinical data were obtained at interview, through case notes and the self-report Eating and Exercise Examination (EEE) which contains the Quality of Life for eating disorders (QOL ED) Global score and examines ED behaviours (objective binge eating, excessive exercise, laxative abuse, self-induced vomiting) over the previous three months (Abraham, Brown, Boyd, Luscombe, & Russell, 2006; Abraham & Lovell, 1999).

2.3. Procedure

Participants were interviewed during admission to a specialised ED unit from 2006 to 2011 (once medically stable and after at least three weeks of regular supervised eating and weight maintenance or weight gain as appropriate). The interviews were conducted by specialists trained in using the IPDE and checked by one expert clinician. Approval was granted by the Human Ethics Committees of the University of Sydney and the Northside Clinic.

2.4. Statistical analysis

A large sample size would be required if all 80 IPDE variables were included in a factor analysis. Consequently, measures were taken to reduce the number of variables included in the factor analysis. Specifically, only items showing adequate response variation, defined as symptom presence of ≥ 25% were retained, keeping the ratio of ten subjects for each question which is common in the literature (Costello & Osborne, 2005) and in keeping with Gorsuch's (1983) recommendation of five subjects per item, with a minimum of 100 subjects. The number of factors was confirmed by scree plot. Principal components analysis using varimax rotation was conducted on the responses to the remaining eleven questions (detailed in Table 1). Sampling adequacy was assessed using the Kaiser–Meyer–Olkin (KMO) statistic and the strength of the relationship amongst the variables with Bartlett's test of sphericity. Factors with an eigenvalue > 1.0 (Comrey & Lee, 1994) and factor loading > 0.50 (Field, 2005) were retained.

Factor scores were calculated by multiplying item scores by the item loading and summing. Univariate ANOVA, controlling for age, compared factor scores by ED behaviours (presence of self-induced vomiting, laxative use, objective binge eating at least once/week, and excessive exercise on more than 19 days/month). Post hoc univariate ANOVA were run for the four ED diagnoses for each factor with age, BMI and purging as covariates. The data were analysed using SPSS version 19.0 and alpha was set at 0.05.

3. Theory/calculation

Researchers have demonstrated that many PD features originally construed as uni-dimensional are in fact not. For example, Lampard et al. (2012) demonstrated two dimensions of perfectionism in ED patients: self-oriented and socially prescribed; and Grilo (2004) reported DSM-IV obsessive-compulsive PD had three factors amongst BED patients: rigidity (interpersonal aspect), perfectionism (intrapersonal aspect) and miserliness (behavioural aspect). These researchers concluded that the results are important in regards to the development, maintenance and treatment of ED comorbid with axis II features, in particular the 'intrapersonal', 'self-oriented' feature of 'perfectionism'

**Table 1**  
Factors arising from factor analysis of the 20 most frequently cited IPDE questions (principal components analysis with varimax rotation).

Factor	Original IPDE Scale	Statement	1	2	3	4	5
Interpersonal anxiety	Avoidant	Inhibited in new interpersonal situation; quieter and more cautious than usual as feels inadequate, unsure, inferior	<b>.838</b>	.074	.043	-.058	.074
	Avoidant	Preoccupation with being criticised or rejected in social situations	<b>.735</b>	-.030	.250	.046	-.022
	Avoidant	Views self as socially inept, unappealing, inferior	<b>.705</b>	.356	.072	-.019	.093
	Avoidant	Reluctant to take personal risks or engage in new activities as may prove embarrassing	<b>.612</b>	.213	.209	.182	.066
	Avoidant	Not willing to get involved with people unless certain of being liked	<b>.591</b>	-.016	-.012	.263	.022
Instability	Borderline	Chronic feelings of emptiness	.136	<b>.803</b>	.113	-.088	.239
	Borderline	Recurrent suicidal behaviour, gestures, threats or self-mutilating behaviour	.152	<b>.766</b>	.168	.065	.025
Self-uncertainty	Borderline	Affective instability due to reactivity or mood	-.003	.486	.290	.532	-.297
	Histrionic	Suggestible/easily influenced by others/ circumstances which causes stress	.046	.112	<b>.774</b>	-.138	.212
	Dependent	Needs excessive amount of advice and reassurance from others for everyday decision	.153	.142	<b>.736</b>	.193	-.022
	Borderline	Identity disturbance/unstable self-image – uncertain about expectancy of own behaviour	.359	.206	<b>.582</b>	.067	-.074
Obsessionality	Obsessive-compulsive	Rigid and stubborn	.213	-.031	-.115	<b>.765</b>	-.003
	Obsessive-compulsive	Excessive devotion to work to exclusion of leisure activities and friends	.081	.028	.091	<b>.706</b>	.294
	Obsessive-compulsive	Reluctant to delegate tasks unless they are done his/her way	-.023	.048	.258	<b>.537</b>	.391
Perfectionism	Obsessive-compulsive	Perfectionism interferes with task completion	.196	.066	-.090	.101	<b>.813</b>
	Obsessive-compulsive	Preoccupied with details, rules, lists, order, schedules, so point of activity is lost	-.001	.069	.196	.248	<b>.708</b>

(Lilenfeld et al., 2000; Shafran, Cooper, & Fairburn, 2002). Our aim was therefore to investigate the factor structure of the IPDE in a sample of patients with ED.

#### 4. Results

Factor analysis yielded five factors accounting for 62% of the total variance (confirmed by scree plot), named 'interpersonal anxiety', 'instability', 'self-uncertainty', 'obsessionality' and 'perfectionism'. The relationship amongst the variables was strong (Bartlett's Test of Sphericity  $\chi^2 = 625.38$ ,  $df = 120$ ,  $p < 0.001$ ), and sampling was adequate (KMO = 0.77). Factors and their original PD classification (DSM-IV) are given in Table 1.

Univariate analysis indicated significant differences on some factors for BMI <18.5 kg/m<sup>2</sup> and purging (self-induced vomiting or laxative use) groups but not for excessive exercise or binge eating (see Table 2). Age was a significant covariate for 'instability' ( $F = 4.81$ ,  $p < 0.03$ ) and 'self-uncertainty' ( $F = 4.22$ ,  $p < 0.04$ ) with less PD as age increases. There were no significant differences in factor scores between ED diagnostic groups when BMI, purging behaviour and age were entered as covariates.

#### 5. Discussion

This study is the first to use items of the IPDE to investigate personality in ED patients using factor analysis. Five factors emerged; four were similar to factors found previously (Goldner, Srikameswaran, Schroeder, Livesley, & Birmingham, 1999; Thompson-Brenner, Eddy, Satir, Boisseau, & Westen, 2008). The new factor 'self-uncertainty' comprised three items, each from a different Axis II diagnosis: being stressed by being easily influenced by others (histrionic), needing excessive amounts of reassurance for everyday decisions (dependent) and an unstable self-image characterised by not knowing what to expect of self in different situations (borderline). Previous factor-analytical studies of personality pathology in ED patients have not found a factor describing this poor self-concept, insecurity and lack of confidence observed in many ED patients (Petrie, Greenleaf, Reel, & Carter, 2009).

The factor 'interpersonal anxiety' contained only items from the IPDE 'avoidance' scale. An 'avoidance' factor has been extracted previously (Espelage, Mazzeo, Sherman, & Thompson, 2002; Goldner et al., 1999; Thompson-Brenner et al., 2008). In this study an association with purging behaviour and lower BMI were found. Whether greater 'interpersonal anxiety' is causal, resulting from or reinforcing maintenance of lower weight and purging behaviour, is unclear.

As with previous studies, a 'perfectionism' factor was observed distinct from 'obsessionality' (Grilo, 2004; Lampard et al., 2012; Thompson-Brenner et al., 2008; Westen & Harnden-Fischer, 2001). Both factors contained items from the IPDE 'obsessive-compulsive' scale, the former emphasising interference with task completion and the latter features like non-acceptance of help, excessive devotion to

work and rigidity. 'Fearing loss of control' and 'preoccupation with eating and weight controlling thoughts' differentiate ED from non-ED (Abraham, von Lojewski, Anderson, Clarke, & Russell, 2009) and may explain the division of the 'obsessive-compulsive' scale of the IPDE into two factors. Whether this would occur in a different population will need investigation.

Similar to many previous studies (Goldner et al., 1999; Thompson-Brenner et al., 2008; Westen & Harnden-Fischer, 2001) the factor 'instability' included features like impulsivity and self-harming behaviours, consisted only of items from the IPDE borderline scale and is highly significantly related to purging behaviour. Greater psychopathology amongst purging ED patients is well recognised (Beumont, George, & Smart, 1976). Purging may account for studies showing differences in PD between different ED diagnostic groups, with borderline features usually BN and binge/purge patients (Claes et al., 2006; Goldner et al., 1999; Thompson-Brenner et al., 2008). The association of 'instability' and 'self-uncertainty' with age was weak but consistent with improvement of PD in ED patients over time and with remission (Klump et al., 2004). This emphasises the need to address self-esteem when treating EDs.

The strengths of this study include: the use of an examination that allows clarification of responses to ensure separation of PD from ED, and using inpatients to ensure that patients were medically stable. The limitations included: sample size restrictions and not having a comparison untreated ED group.

The personality factors found in this study include those cited as contributing to the chronicity of EDs. Treasure et al. (2011) hypothesised that "negative schema common to ED, for example low self-esteem, perfectionism and striving for social value may augment existing or initiate new eating disorder behaviour" (p. 296). These factors, which may contribute to chronicity should be managed early in treatment. The personality factors found may also suggest further assessment of early attachment may improve treatment and outcome.

#### Role of funding source

There were no involvements of funding resources in this project.

#### Contributors

Astrid von Lojewski and Suzanne Abraham developed the idea of the research, analysed the data with the advice of Dr Georgina Luscombe and discussed the results. Astrid von Lojewski did the literature research and wrote the article. Suzanne Abraham coordinated the data collection and proof-read the article. Both authors contributed to and have approved the final manuscript.

#### Conflict of interest

Both authors declare that there is no conflict of interest.

#### Acknowledgements

We thank Georgina Luscombe for her statistical advice.

**Table 2**

Mean factor loading for eating disorder patients' behaviours<sup>a</sup> and BMI in the previous three months<sup>b</sup>: purging and a BMI below 18.5 kg/m<sup>2</sup>.

	Purging (vomit/laxative)		F	p	BMI less than 18.5 kg/m <sup>2</sup>		F	p
	Present	Absent			Yes	No		
	N = 102	N = 53			N = 75	N = 80		
	Mean (SD)	Mean (SD)			Mean (SD)	Mean (SD)		
Interpersonal anxiety	2.5 (2.1)	2.1 (2.2)	4.85	0.029	2.6 (2.1)	2.1 (2.0)	7.48	0.007
Instability	1.8 (1.4)	0.6 (0.9)	17.65	0.000	1.1 (1.2)	1.8 (1.5)	3.65	ns
Self-uncertainty	0.8 (1.1)	0.5 (0.9)	3.50	ns	0.8 (1.1)	0.7 (1.0)	0.76	ns
Obsessionality	0.9 (1.0)	0.9 (1.1)	0.42	ns	1.0 (1.2)	0.8 (0.9)	3.24	ns
Perfectionism	0.9 (1.0)	0.5 (0.8)	9.32	0.003	0.8 (1.0)	0.7 (0.9)	3.00	ns

<sup>a</sup> No significant differences for the behaviours of binge eating and excessive exercise for any factors.

<sup>b</sup> Results of univariate ANOVA with age as a covariate; age significant for 'instability'  $F = 4.81$ ,  $p < 0.03$  and 'self-uncertainty'  $F = 4.22$ ,  $p < 0.04$ .

## References

- Abraham, S. F., Brown, T., Boyd, C., Luscombe, G., & Russell, J. (2006). Quality of life: Eating disorders. *Australian and New Zealand Journal of Psychiatry*, 40, 150–155.
- Abraham, S. F., & Lovell, N. (1999). Research and clinical assessment of eating and exercise behaviour. *Hospital Medicine*, 60, 481–485.
- Abraham, S., von Lojewski, A., Anderson, G., Clarke, S., & Russell, J. (2009). Feelings: What questions best discriminate women with and without eating disorders? *Eating and Weight Disorders*, 14, 6–10.
- American Psychiatric Association (APA) (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington DC: Author.
- Beumont, P. J., George, G. C., & Smart, D. E. (1976). "Dieters" and "vomitters and purgers" in anorexia nervosa. *Psychological Medicine*, 6, 617–622.
- Cassin, S. E., & von Ranson, K. M. (2005). Personality and eating disorders: A decade in review. *Clinical Psychological Review*, 25, 895–916.
- Claes, L., Vandereycken, W., Luyten, P., Soenens, B., Pieters, G., & Vertommen, H. (2006). Personality prototypes in eating disorders based on the Big Five model. *Journal of Personality Disorders*, 20, 401–416.
- Comrey, A. L., & Lee, H. B. (1994). *A first course in factor analysis*. Hillsdale, NJ: Lawrence Erlbaum Associates Inc.
- Costello, A.B., & Osborne, J. (2005). Best practices in exploratory factor analysis: Four recommendations for getting the most from your analysis. *Practical Assessment, Research & Evaluation*, 10 (Retrieved January 28, 2013, from <http://pareonline.net/pdf/v10n7.pdf>).
- Diaz-Marsa, M., Carrasco Perera, J. L., Prieto Lopez, R., & Saiz Ruiz, J. (2000). Role of personality in eating behaviour disorders. *Actas Españolas de Psiquiatría*, 28, 29–36.
- Espelage, D. L., Mazzeo, S. E., Sherman, R., & Thompson, R. (2002). MCMI-II profiles of women with eating disorders: A cluster analytic investigation. *Journal of Personality Disorders*, 16, 453–463.
- Field, A. (2005). *Discovering statistics using SPSS*. Singapore: Sage Publications Ltd.
- Godt, K. (2008). Personality disorders in 545 patients with eating disorders. *European Eating Disorders Review*, 16, 94–99.
- Goldner, E. M., Srikaneswaran, S., Schroeder, M. L., Livesley, W. J., & Birmingham, C. L. (1999). Dimensional assessment of personality pathology in patients with eating disorders. *Psychiatry Research*, 85, 151–159.
- Gorsuch, R. L. (1983). *Factor analysis*. Hillsdale, NJ: Erlbaum.
- Grilo, C. M. (2002). Recent research of relationships among eating disorders and personality disorders. *Current Psychiatry Reports*, 4, 18–24.
- Grilo, C. M. (2004). Factor structure of DSM-IV criteria for obsessive compulsive personality disorder in patients with binge eating disorder. *Acta Psychiatrica Scandinavica*, 109, 64–69.
- Haslem, N., Holland, E., & Kuppens, P. (2012). Categories versus dimensions in personality and psychopathology: A quantitative review of taxometric research. *Psychological Medicine*, 42, 903–920.
- Ilkjaer, K., Kortegaard, L., Hoerder, K., Joergensen, J., Kyvik, K., & Gillberg, C. (2004). Personality disorders in a total population twin cohort with eating disorders. *Comprehensive Psychiatry*, 45, 261–267.
- Karwautz, A., Troop, N. A., Rabe-Hesketh, S., Collier, D. A., & Treasure, J. L. (2003). Personality disorders and personality dimensions in anorexia nervosa. *Journal of Personality Disorders*, 17, 73–85.
- Kennedy, S. H., McVey, G., & Katz, R. (1990). Personality disorders in anorexia nervosa and bulimia nervosa. *Journal of Psychiatric Research*, 24, 259–269.
- Klump, K. L., Strober, M., Bulik, C. M., Thornton, L., Johnson, C., Devlin, B., et al. (2004). Personality characteristics of women before and after recovery from an eating disorder. *Psychological Medicine*, 34, 1407–1418.
- Lampard, A.M., Byrne, S. M., McLean, N., & Fursland, A. (2012). The Eating Disorder Inventory-2 Perfectionism scale: Factor structure and associations with dietary restraint and weight and shape concern in eating disorders. *Eating Behaviors*, 13, 49–53.
- Larsson, J. O., & Hellzen, M. (2004). Patterns of personality disorders in women with chronic eating disorders. *Eating and Weight Disorders*, 9, 200–205.
- Lilenfeld, L. R., Stein, D., Bulik, C. M., Strober, M., Plotnicov, K., Pollice, C., et al. (2000). Personality traits among currently eating disordered, recovered, and never ill first-degree female relatives of bulimic and control women. *Psychological Medicine*, 30, 1399–1410.
- Lilenfeld, L. R., Wonderlich, S., Riso, L. P., Crosby, R., & Mitchell, J. (2006). Eating disorders and personality: A methodological and empirical review. *Clinical Psychological Review*, 2, 299–320.
- Loranger, A. W., Janca, A., & Sartorius, N. (1997). *International Personality Disorders Examination (IPDE)*. Geneva: World Health Organization.
- Loranger, A. W., Sartorius, N., Andreoli, A., Berger, P., Buchheim, P., Channabassavanna, S. M., et al. (1994). The International Personality Disorder Examination: The World Health Organisation/alcohol, drug abuse and mental health administration international pilot study of personality disorders. *Archives General Psychiatry*, 51, 125–224.
- Milos, G. F., Spindler, A., Buddeberg, C., & Cramer, A. (2003). Axes I and II comorbidity and treatment experiences in eating disorder subjects. *Psychotherapy and Psychosomatics*, 72, 276–285.
- Modestin, J., Oberson, B., & Erni, T. (1997). Possible correlates of DSM-III-R personality disorders. *Acta Psychiatrica Scandinavica*, 96, 424–430.
- Petrie, T. A., Greenleaf, C., Reel, J., & Carter, J. (2009). Personality and psychological factors as predictors of disordered eating among female collegiate athletes. *Brunner-Mazel Eating Disorder Monograph Series*, 17, 302–321.
- Rosenvinge, J. H., Martinussen, M., & Ostensen, E. (2000). The comorbidity of eating disorders and personality disorders: A meta-analytic review of studies published between 1983 and 1998. *Eating and Weight Disorders*, 5, 52–61.
- Shafraan, R., Cooper, Z., & Fairburn, C. G. (2002). Clinical perfectionism: A cognitive-behavioral analysis. *Behaviour Research and Therapy*, 40, 773–791.
- Skodol, A. E., Oldham, J. M., Hyler, S. E., Kellman, H. D., Doidge, N., & Davies, M. (1993). Comorbidity of DSM-III-R eating disorders and personality disorders. *International Journal of Eating Disorders*, 14, 403–416.
- Steiger, H., & Bruce, K. (2004). Personality traits and disorders in anorexia nervosa, bulimia nervosa, and binge eating disorder. In T. D. Brewerton (Ed.), *Clinical handbook of eating disorders: An integrated approach* (pp. 209–230). New York, NY: Marcel Dekker Inc.
- Thompson-Brenner, H., Eddy, K. T., Satir, D. A., Boisseau, C. L., & Westen, D. (2008). Personality subtypes in adolescents with eating disorders: Validation of a classification approach. *Journal of Child Psychology and Psychiatry*, 49, 170–180.
- Treasure, J., Crane, A., McKnight, R., Buchanan, E., & Wolfe, M. (2011). First do no harm: Iatrogenic maintaining factors in anorexia nervosa. *European Eating Disorders Review*, 19, 296–302.
- van Hanswijck de Jonge, P., van Furth, E. F., Lacey, H., & Waller, G. (2003). The prevalence of DSM-IV personality pathology among individuals with bulimia nervosa, binge eating disorder and obesity. *Psychological Medicine*, 33, 1311–1317.
- Vitousek, K., & Manke, F. (1994). Personality variables and disorders in anorexia nervosa and bulimia nervosa. *Journal of Abnormal Psychology*, 103, 137–147.
- Wagner, A., Barbarich-Marsteller, N. C., Frank, G. K., Bailer, U. F., Wonderlich, S. A., Crosby, R. D., et al. (2006). Personality traits after recovery from eating disorders: Do subtypes differ? *International Journal of Eating Disorders*, 39, 276–284.
- Westen, D., & Harnden-Fischer, J. (2001). Personality profiles in eating disorders: Rethinking the distinction between axis I and axis II. *American Journal of Psychiatry*, 158, 547–562.
- Wonderlich, S. A., Crosby, R. D., Engel, S. G., Mitchell, J. E., Smyth, J., & Miltenberger, R. (2007). Personality-based clusters in bulimia nervosa: Differences in clinical variables and ecological momentary assessment. *Journal of Personality Disorders*, 21, 340–357.
- Wonderlich, S., & Mitchell, J. E. (2001). The role of personality in the onset of eating disorders and treatment implications. *Psychiatry Clinics of North America*, 24, 221–225.
- Zubieta, J. K., Demitrack, M.A., Fenick, A., & Krahn, D.D. (1995). Obsessionality in eating-disorder patients: Relationship to clinical representation and two-year outcome. *Journal of Psychiatric Research*, 29, 333–342.